

**Thermodynamics for Biological Systems:
Classical and Statistical Aspects
Prof. G.K. Suraishkumar
Department of Biotechnology
Indian Institute of Technology - Madras**

**Lecture – 44
Liquid-Liquid and Solid-Liquid Equilibria**


Welcome!

In the last class, we had looked at vapour-liquid equilibrium as a special case of equilibrium; and we had worked out the special condition for vapour-liquid equilibrium based on our original conditions of equilibrium – the five equations that we had seen very early in this module. And we had also worked out an example problem to better understand how the equilibrium comes about, and how we can use the various parameters that we have developed or defined to come about with useful relationships at equilibrium, in the case of vapour-liquid equilibrium.

What we are going to do in this class, essentially, is to develop the governing equations for liquid-liquid and solid-liquid equilibria. There could be various different kinds of equilibria. We are just going to take these two, and develop the governing equations, because these occur quite often in the case of biological systems, especially bio-process related activities as well as some activities that deal with analysis of biological molecules and so on.

What we are going to do, how we are going to do develop these governing equations is as exercises, which means you are going to develop it. That way, it builds up your confidence of addressing and developing equations in hitherto unknown situations. A complete appreciation of the principles that are involved also comes about as a part of doing such exercises. So, we will do these as exercises. I am going to start you out, then I will give time, and of course, I will present the equations, the governing equations, the way I have done it. We can cross check after sometime.

(Refer Slide Time: 02:26)



What I would like you to remember, when you are developing these governing equations for liquid-liquid and solid-liquid equilibria ... is the fact that equations 5.1 to 5.5 and further – you know we had stopped at 5.5. Essentially equation 5.1 is

$$T_{\alpha} = T_{\beta} = T_{\gamma} = \dots$$

the sameness of temperature or the equality of temperature across the various phases at equilibrium. The equation 5.2 was the equation of pressures or equality of pressures across the various phases,

$$P_{\alpha} = P_{\beta} = P_{\gamma} = \dots$$

Then equations 5.3 to 5.5, we had written for three components; one, two and three. What we had written there was that the chemical potential of each component was equal across the various phases, in which it is present, of course. We are assuming that all the components are present in all the phases. Therefore

$$\mu_{1,\alpha} = \mu_{1,\beta} = \mu_{1,\gamma} = \dots$$

was equation 5.3.


$$\mu_{2,\alpha} = \mu_{2,\beta} = \mu_{2,\gamma} = \dots$$

equation 5.4.

$$\mu_{3,\alpha} = \mu_{3,\beta} = \mu_{3,\gamma} = \dots$$

was equation 5.5. These are the 5 fundamental equations that are necessary for the thermodynamic equilibrium in any system. Need to keep this in mind, and we need to develop it.

(Refer Slide Time: 04:19)




Liquid-Liquid equilibrium

The distinct liquid phases in equilibrium could be of different kinds

- one hydrophilic and the other hydrophobic, as in the case of antibiotic extraction process
- or two aqueous phases, as in the case of aqueous two phase purification of proteins

If /1 and /2 are the two liquid phases in equilibrium, for each component, i , in the two phases, we can write

 NPTEL

Let me start you out for the case of liquid-liquid equilibrium, which means there are two liquid phases ... distinctly separate. ... They could be of many different kinds. One could be a hydrophilic phase, the other could be hydrophobic phase. It is easy to imagine a hydrophilic phase – hydrophobic phase; let us say, a mixture of oil and water. They are not going to mix with each other. You can clearly demarcate these two phases. One such system may be toluene-water system and so on, so forth, is actually used in the anti-biotic extraction process. One hydrophilic phase, which is the water phase, in which the anti-biotic is present after it is produced by the organisms, after the bio-reactor phase. And, what is added may be hydrophobic liquid is added to extract most of the anti-biotic into the hydrophobic phase; because, at equilibrium, the concentration of the anti-biotic in the hydrophobic phase is many fold the concentration of the anti-biotic in the hydrophilic or the water phase. That is the basic reason for using this two liquid process for extraction and purification of certain bio- molecules such as antibiotics. So, this is

one example of a liquid-liquid ... or of the occurrence of two liquid phases. ... Equilibrium conditions are the limiting conditions. More interestingly, they could be even two aqueous phases, two water based phases, but which are distinct phases ... one sits on the top of the other.

An example is different mixtures of let us say polyethylene glycol and ammonium sulphate salt. If we mix them together then depending on the composition they could form two different, distinct phases. In fact such two aqueous phase system, or aqueous two phase systems as they are called, are actually used in the industry for extraction and purification of proteins. Why do we need to do that?

You know that proteins get their activity because of the way the amino acid chain is folded. And the way the amino acid chain is folded ... has relevance only in an aqueous system, because, if you recall from your early classes, may be an initial life science course or molecular biology course, biochemistry course for that matter ... that a protein folds itself according to the forces between them as well as the forces between the various constituents of the proteins, their amino acids, their side chains and so on, and the environment that it is present in.

If the environment that it is present in is water, it forms a certain conformation which is active. You put the same protein molecule in a non-aqueous phase then the forces are completely different. And the way it gets folded may result in a molecule that is not active at all; because, it is only the three dimensional folding of the protein molecule that gives that its activity. Therefore, what is done is, two aqueous phases are used. And that is based on the fact that the solubility of protein in one of the aqueous phases is higher may not be 10 or 100 fold higher but may be about a few fold ... 3 to 5 fold higher than the other phase. And this was used to concentrate and purify the protein molecules.

Therefore, liquid-liquid systems are used quite extensively in biological systems, and we are going to look at ... liquid-liquid systems, in general. What I would like you to do is take about 20 minutes from now; because you need to think about various things and so on. And the question is this, if l_1 and l_2 are the two liquid phases in equilibrium, for each component i in the two phases, develop the expression that can be written. It is as simple as that. To give you a hint, the first five equations, equations 5.1 to 5.5 are valid. Also take a look at how we went about doing pretty much the same thing, but for a component i , that is distributing between the vapour and liquid phases in the case of vapour-liquid equilibrium. Those are the hints that I am going to leave you with. Take about 20 minutes, and then we will look at what the solution is.

(Refer Slide Time: 09:20)

Please pause the video here. You can pause it as long as you want. The time mentioned is only a guideline

So, you would have found it easy to write the condition for equilibrium which is nothing but the equality of chemical potentials. We are of course, going to take the conditions of temperature equality and pressure equality as a given. So we do not have to write them explicitly here as mentioned some time earlier; that is considered as a given thing. Here

$$\mu_i^{l1} = \mu_i^{l2}$$

i is the component which is distributing between the two liquid phases $l1$ and $l2$. As mentioned earlier $l1$ could be a hydrophilic phase and $l2$ could be a hydrophobic phase. Or $l1$ could be one aqueous phase and $l2$ could be another aqueous phase, as in the case of aqueous two phase extraction. Let us call this equation 5.39.

(Refer Slide Time: 10:40)


Representing Eq. 5.39 in terms of fugacities, we get

$$\mu_i^0 + RT \ln \hat{f}_i^{l1} = \mu_i^0 + RT \ln \hat{f}_i^{l2}$$

And thus

$$\hat{f}_i^{l1} = \hat{f}_i^{l2} \quad \text{Eq. 5.40}$$

Using Eq. 4.8, we get

$$\gamma_i^{l1} f_i^{l1} x_i^{l1} = \gamma_i^{l2} f_i^{l2} x_i^{l2} \quad \text{Eq. 5.41}$$


Now, if we expand equation 5.39 in terms of fugacities, we can write

$$\mu_i^0 + RT \ln \hat{f}_i^{l1} = \mu_i^0 + RT \ln \hat{f}_i^{l2}$$

This is the fugacity of component i in $l1$ in a multi-component system. And similarly, this \hat{f}_i^{l2} is the fugacity of component i in $l2$, in a multi-component system. Therefore, it is quite easy to see ... μ_i^0 is going to cancel with μ_i^0 ; RT cancels with this side ... this side. And then $\ln \hat{f}_i^{l1}$ equals $\ln \hat{f}_i^{l2}$. And therefore,

$$\hat{f}_i^{l1} = \hat{f}_i^{l2}$$

Now if you go back and look at expressions for this, you could write ... which is essentially equation 4.8 in the previous module. You could express \hat{f}_i^{l1} in terms of the pure component fugacity, which is easier to estimate. This is written as

$$\gamma_i^{l1} f_i^{l1} x_i^{l1} = \gamma_i^{l2} f_i^{l2} x_i^{l2}$$

This becomes the condition for equilibrium. Let us call this equation 5.41.

(Refer Slide Time: 12:48)

Example 5.4

For the liquid-liquid extraction of ampicillin, at equilibrium, express the activity coefficients of ampicillin in the two liquid phases in terms of fugacities and mole fractions

We know from Eq. 5.39 that

$$\mu_{amp}^{l1} = \mu_{amp}^{l2}$$

$$\gamma_{amp}^{l1} f_{amp}^{l1} x_{amp}^{l1} = \gamma_{amp}^{l2} f_{amp}^{l2} x_{amp}^{l2}$$

Thus

$$\frac{\gamma_{amp}^{l1}}{\gamma_{amp}^{l2}} = \frac{f_{amp}^{l2} x_{amp}^{l2}}{f_{amp}^{l1} x_{amp}^{l1}}$$



To get a little more comfortable with this, let us work out an example. For the liquid-liquid extraction of ampicillin, at equilibrium, express the ratio of the activity coefficients of ampicillin – that ratio is missing here – the ratio of activity coefficients of the ampicillin in the two liquid phases in terms of fugacities and mole fractions. Take about 15 minutes. Go ahead and do this please.

This is on the same lines as the derivation. We could directly use the final relationship that we got just a while ago. We know, if we employ equation 5.39 directly, ... see, this is the case of liquid-liquid equilibrium and ampicillin is the component *i* that is distributing between the two liquid phases. We know that from equation 5.39 that

$$\mu_{amp}^{l1} = \mu_{amp}^{l2}$$

Writing this in terms of gamma *i* or bringing this down to fugacities, and then writing it; either way it is ok.


$$\gamma_{amp}^{l1} f_{amp}^{l1} x_{amp}^{l1} = \gamma_{amp}^{l2} f_{amp}^{l2} x_{amp}^{l2}$$

Therefore, what is required is the ratio of the activity coefficients of ampicillin,

$$\frac{\gamma_{amp}^{l1}}{\gamma_{amp}^{l2}} = \frac{f_{amp}^{l2} x_{amp}^{l2}}{f_{amp}^{l1} x_{amp}^{l1}}$$

This is the equation that we were looking for, or the relationship that we are looking for in this example.

(Refer Slide Time: 15:12)




Solid-Liquid equilibrium

Solid-liquid equilibria arise in biological situations such as biosorption, bioleaching, and even in some analytical steps

Using similar arguments as for the liquid-liquid equilibrium, we can start with the governing equation

$$\mu_i^s = \mu_i^l \quad \text{Eq. 5.42}$$

 NPTEL

For completeness, I would ... let me actually present this since we are almost out of time in this particular lecture. Let me just present this. It is quite straight forward. It is an example of solid-liquid equilibrium. There are many different examples of solid-liquid equilibrium in biological systems. They arise in situations such as biosorption which is the removal of, let us say, toxic trace metals using a biological agents such as cells and so on. Bioleaching which is taking the metals or useful components out by using biological means, and even in some analytical steps you come across solid-liquid equilibrium in biological systems. ... We know that equations 5.1 to 5.5 are valid. Therefore, using similar arguments as for the liquid-liquid equilibrium, we can start with the governing equation that

$$\mu_i^s = \mu_i^l$$

Equation 5.42.

(Refer Slide Time: 16:29)

Thus

$$\hat{f}_i^s = \hat{f}_i^l$$

and, in terms of activity coefficients, we can write

$$\gamma_i^s f_i^s z_i = \gamma_i^l f_i^l x_i$$



Therefore,

$$\hat{f}_i^s = \hat{f}_i^l$$

In terms of activity coefficient,

$$\gamma_i^s f_i^s z_i = \gamma_i^l f_i^l x_i$$

the mole fraction in the solid phase is given as z_i . This is the governing equation for solid-liquid equilibrium. When we come back in the next class, we will review whatever we have done in this particular module on phase equilibrium. See you then.