


Introduction to Nonlinear Dynamics
Prof. Gaurav Raina
Department of Electrical Engineering
Indian Institute of Technology, Madras

Module -04
Lecture-10
1-Dimensional Flows, Bifurcations, Lecture 2

(Refer Slide Time: 00:01)




The first bifurcation is the saddle node bifurcation. This is the basic mechanism by which fixed points are either created or destroyed, essentially as a system parameter is varied. The two fixed points move towards each other, collide and then mutually annihilate. Prototypical example is $\dot{x} = r + x^2$ where r is a parameter which is greater than 0, is equal to 0 or less than 0. So, for r less than 0, if you plot \dot{x} versus x , that is the plot for \dot{x} versus x , we find that we have two fixed points.

So, the system has two fixed points one is stable and the other is unstable, for r is equal to 0. When we plot \dot{x} versus x , we find that we only have one fixed point. The fixed point is attracting from the left and repelling from the right. So, the fixed points actually turn into a half-stable fixed point at $x^* = 0$ and when r is greater than 0 plotting \dot{x} versus x reveals that in fact we have no fixed points.

So, a bifurcation effectively takes place at $r = 0$ as the vector fields for r greater than 0 and r less than 0 are qualitatively different from each other.

(Refer Slide Time: 02:11)



Now, we go ahead and plot something known as bifurcation diagrams. So, if we show a stack of vector fields for discrete values of r that is the stack of vector fields, when r greater than 0 there are no fixed points. When r is equal to 0, there is half stable fixed point and when r less than 0, we have two fixed points, one stable and other unstable and that is what happens when r varies, so we have stable and unstable branches.

So, a common way is to treat the parameter r as playing the role of an independent variable in the limit of a continuous stack of vector fields. What we get is the following, so we plot x versus r as the stable branch and that is the unstable branch. So, we can also include arrows in the diagram, so including arrows in the bifurcation diagram we get. So, this is called the bifurcation diagram for the saddle node bifurcation. So, this is the first bifurcation that we have dealt with and diagram on the left is referred to as its bifurcation diagram.

(Refer Slide Time: 04:15)

We now conduct a linear stability analysis of $\dot{x} = r - x^2$

First identify the fixed points $\dot{x} = f(x) = r - x^2$ gives $x^* = \pm\sqrt{r}$

For $r > 0$, we get two fixed points

To establish linear stability, we get $f'(x^*) = -2x^*$

So $x^* = +\sqrt{r}$ is stable
 $x^* = -\sqrt{r}$ is unstable


For $r < 0$, there are no fixed points

For $r = 0$, we get $f'(x^*) = 0$

The linearized term vanishes!

Let us now conduct a linear stability analysis of $\dot{x} = r - x^2$, we first identify the fixed points, so $\dot{x} = f(x) = r - x^2$, yields (gives) $x^* = \pm \sqrt{r}$. For r greater than 0, we get two fixed points and to establish linear stability, we get $f'(x^*) = -2x^*$. So $x^* = +\sqrt{r}$ yields a stable fixed point, $x^* = -\sqrt{r}$ is unstable. For r less than 0, there are actually no fixed points, for $r = 0$, we get $f'(x^*) = 0$ and so the linearized term actually vanishes.

(Refer Slide Time: 05:25)



Let us consider an example, consider the first order system $\dot{x} = r - x - e^{-x}$, show that it undergoes a saddle node bifurcation as r varies, further finding the value of r at the bifurcation point. First, we plot $r - x$ and e^{-x} on the same graph, where the line $r - x$ intersects with the

curve e^{-x} to the $-x$, we get $r - x = e^{-x}$ to the $-x$ and so $f(x)$ which is $r - x - e^{-x} = 0$. So, intersections of the line and the curve actually correspond to the fixed points of the system.

Now let us go ahead and make some plots, that is the line $r - x$ that is the curve e^{-x} to the $-x$. So we note that this system has two fixed points, one stable and the other one is unstable. Note that the flow is to the right where $r - x$ is greater than e^{-x} and so \dot{x} is greater than 0. Now let us go ahead and decrease the parameter r little bit, so that is the line $r - x$, that is curve e^{-x} to the $-x$ and note that in this case we have one fixed point.

The fixed point turns out to be actually a half stable fixed point. So, at some critical value $r = r_c$ the line actually becomes tangent to the curve. So that is the saddle node bifurcation point, for r less than r_c that is $r - x$ that e^{-x} to the $-x$ and note that they do not actually intersect at all, so there are no fixed points.

(Refer Slide Time: 08:05)

We still need to find the bifurcation point r_c

Impose the condition that the graphs of $r-x$ and e^{-x} intersect tangentially

We require equality of the functions and their derivatives

So $e^{-x} = r-x$

and $\frac{d}{dx} e^{-x} = \frac{d}{dx} (r-x)$

$\Rightarrow -e^{-x} = -1$

$\Rightarrow x = 0$

Substituting $x=0$ into $e^{-x} = r-x$ gives $r=1$

So the bifurcation point $r_c = 1$ and the bifurcation occurs at $x=0$

We still need to find the bifurcation point r_c . So, we impose the condition that the graphs of $r - x$ and e^{-x} intersect tangentially. We require equality of the functions and their derivatives. So, $e^{-x} = r - x$ and $\frac{d}{dx} e^{-x} = \frac{d}{dx} (r - x)$ which gives us $-e^{-x} = -1$, which yields $x = 0$, so substituting $x = 0$ into $e^{-x} = r - x$ gives $r = 1$. So, the bifurcation point is at $r_{\text{critical}} = 1$ and the bifurcation occurs at $x = 0$.

(Refer Slide Time: 09:19)



In this lecture, we introduced the saddle node bifurcation. This is the basic mechanism through which fixed points can be created or destroyed. We introduced prototypical examples for the saddle node. Now essentially what it showed us was the following, we had a model, where we actually had two fixed points, one of the fixed point was stable and one of the fixed points was unstable. And then a parameter in the system changes as the parameter changes, both of these fixed points actually came close to each other.

And at particular point of time they actually merged and became one half stable fixed point and as the parameter change even more, in fact the fixed point in the system vanished. So, we had a scenario, where we had two fixed points stable and an unstable. Then we had one fixed point which was half stable fixed point and as the parameter varied all fixed points in the system actually vanished.

So that is why fascinating because when you modelling the real world, you have the real world of which you abstract about a simplified model and that model will have some parameters in it. And so the lesson here is that as parameters varies you would have fairly serious qualitative changes in the underline dynamics and we illustrated that through the saddle node bifurcation in this lecture.