Introduction to Probability Theory and Stochastic Processes Prof. S. Dharmaraja Department of Mathematics Indian Institute of Technology, Delhi


Lecture - 74

(Refer Slide Time: 00:01)

Waiting Time Distribution Consider a time-homogenous CTMC. Al t= 0, × (0)= i h known. Let T be a random variable denoting time taken for a change at state from state i. P[2 > ++ E/× (a)=i] = P[z > +++/x(x)=1)[z>>/x(0)=i] $F_{\tau}^{c}(A+t) = F_{\tau}^{c}(t) F_{\tau}^{c}(A)$

So, before going to the P ij, you see the sample path of the sample path of a time homogeneous continuous time Markov chain.

(Refer Slide Time: 00:10)




As I said the system is a staying for some positive amount of time in any state before moving into any other states. Our interest is what is the distribution or what is the waiting time distribution of system being in any state before moving into any other states, that is our interest to find out.

So, how you are going to find out that I am going to explain. That is called the waiting time distribution; that means, what is the distribution of time spending in any state for a time homogeneous continuous time Markov chain before moving into any other states. I assumed that a time 0 the system was in the state i; that means, X of 0 is equal to i that is known or the probability of X of 0 is equal to i, that probability is 1.

Let me make out the random variable tau, that is a random variable denoting the time taken for a change of state from the state i. Change of state means it does not matter which state it goes my interest is to find out what is the waiting time distribution for the state i. The time spent in the state i for that let me make a simple graph so this is t and this is X of t.

(Refer Slide Time: 01:55)



Suppose you assume that the system was in the state i at the time point 0. After some time, it moved into some other state, at the time point s, it was in the state i. At the time

point t also it moved into some other state. So, the tau here is nothing but the time spent in the state i from here to here so, that is a random variable.

So, what I am going to do? I am going to find out what is the complement a CDF for the random variable tau, that is a what is the probability that the tau greater than s plus t given that X of 0 is equal to i. That is same as the probability of the tau is greater than s plus t, given that I can introduce one more condition tau is greater than s. Then I can multiply by using the total theorem of probability tau greater than s given that X of 0 is equal to i.

That is same as the first one I can rewrite as a probability of tau greater than s plus t, given that X of s is equal to i. Because X of 0 is equal to i as well as tau is greater than s, where tau is a time spent in the state i therefore, I can make out X of s is equal to i by combining these 2 concept multiplied by in the probability of tau greater than s given that X of 0 is equal to i, that is same expression here.


Now, the probability of tau greater than s plus t given that X of s is equal to i. That I can rewrite because the this Markov chain is a time homogeneous Markov chain. So, the s to s plus t that is same as the compliment CDF of the random variable tau for the time t. Because it is a s to s plus t since it is Markov chain is a time homogeneous only the length is matters that is the interval of length t. Therefore, this is nothing but the compliment CDF for the random variable tau the time point t multiplied by this is nothing but 0 to s.

So, this is a the compliment CDF of the random variable tau the time point s. Whereas, the left hand side is the compliment CDF for the random variable tau for the time point s plus t. So, what we got the result is the compliment CDF of the unknown random variable tau at the time point s plus t that is same as the product of compliment CDF at the time point s and t. So, this is valid for all s and t greater than 0.

So, we have to find out what is the random variable or what is the distribution going to satisfies this compliment CDF at the time point s plus t is same as the product of compliment CDF at the time point s and t. If any distribution satisfies this compliment a CDF property, then we can find out the distribution for the random variable top.

So, in this derivation we have use the time homogeneous property as well as the total probability rule as well as we have used the Markov property, therefore, it land up the compliment CDF satisfying this equation.

(Refer Slide Time: 06:32)



Now we have to find out what is the distribution going to satisfies this property. So, if you substitute any function with the e power any parameter lambda with the exponential of e power minus lambda t, the previous equation is going to be satisfied. As long as the function is of the form e power minus lambda t, for lambda is greater than 0 and t is greater than 0.

Since the compliment CDF is e power minus lambda t therefore, the CDF of the unknown random variable tau, that is 1 minus e power minus lambda t for t greater than 0 for some lambda. And you know that if the CDF of the random variable is 1 minus e power minus lambda t for t greater than 0 and lambda greater than 0, then that random variable is a exponentially distributed random variable.

So, we can conclude the amount of time or the time taken by the system staying in the state i that time is a exponentially distributed that is a continuous random variable whose distribution is exponential distribution; with the parameter lambda. Even we can specify lambda suffix i that means, it is going to be a function of it depends on the i; that means, if the random variable is going to spend in some state, and that is always exponential

distribution with some parameter lambda and that parameter lambda may depend on the state i.

That means if I go back to the sample path, I can say that the time the system spending in this particular state that is exponentially distributed with some parameter. Then it moved into some other state, the time spending in this state that is also exponentially distributed with some other, it could be some other parameter it depends on that particular state.

Then it moved into the some other state and the time spending in this state, that is also exponentially distributed, and later we can conclude all these the time spending in each state because of it is a Markov property satisfied, the time spent in this state the time spent in this state all are exponentially distributed which is independent of the other.

So, all are going to be mutually independent random variables, then only the Markov property is going to be satisfied; that means, whenever the system is moving from one state to another state, you will have a exponentially distributed time spending in each state, and they are form a mutually independent.

And since the exponential distribution has the memory less property, the system spending in this state, if you just observe at any time t and what is the probability that the system will be for some more time in the same state, given that it was spending already this much time in the state, then that is also exponential distribution. Because of memoryless property of exponentially distribution and which is independent of how much time spend in the same state already.

Therefore the Markov property is going to be satisfied throughout the time, whether the system spending in this state or the other state and so on. So, the Markov property is going to be satisfied for all the time points and the time spending in each state is exponentially distributed. And all the random variable spending in each state all are going to be mutually independent random variables. Now we found out what is the time spending in each state and that is exponential distribution with some parameter. Lambda i and the distribution is same for all i whereas, the value of the parameter lambda maybe depends on the i.